

Document details

< Back to results | 1 of 1

Export Download Print E-mail Save to PDF Add to List More...

Full Text View at Publisher

Journal of High Energy Physics Open Access
Volume 2018, Issue 9, 1 September 2018, Article number 46

Search for dark matter produced in association with a Higgs boson decaying to $\gamma\gamma$ or $\tau^+\tau^-$ at $\sqrt{s}=13$ TeV (Article) (Open Access)

Sirunyan, A.M.^a, Tumasyan, A.^a, Adam, W.^b, Ambroggi, F.^b, Asilar, E.^b, Bergauer, T.^b, Brandstetter, J.^b, Dragicevic, M.^b, Erö, J.^b, Escalante Del Valle, A.^b, Flechl, M.^b, Frühwirth, R.^{b,gu}, Ghete, V.M.^b, Hrubec, J.^b, Jeitler, M.^{b,gu}, Krammer, N.^b, Krätschmer, I.^b, Liko, D.^b, Mädlener, T.^b, Mikulec, I.^b, Rad, N.^b,

View additional authors

^aYerevan Physics Institute, Yerevan, Armenia
^bInstitut für Hochenergiephysik, Wien, Austria
^cInstitute for Nuclear Problems, Minsk, Belarus

View additional affiliations

Abstract

View references (74)

A search for dark matter particles is performed by looking for events with large transverse momentum imbalance and a recoiling Higgs boson decaying to either a pair of photons or a pair of τ leptons. The search is based on proton-proton collision data at a center-of-mass energy of 13 TeV collected at the CERN LHC in 2016 and corresponding to an integrated luminosity of 35.9 fb⁻¹. No significant excess over the expected standard model background is observed. Upper limits at 95% confidence level are presented for the product of the production cross section and branching fraction in the context of two benchmark simplified models. For the Z' -two-Higgs-doublet model (where Z' is a new massive boson mediator) with an intermediate heavy pseudoscalar particle of mass $m_A = 300$ GeV and $m_{DM} = 100$ GeV, the Z' masses from 550 GeV to 1265 GeV are excluded. For a baryonic Z' model, with $m_{DM} = 1$ GeV, Z' masses up to 615 GeV are excluded. Results are also presented for the spin-independent cross section for the dark matter-nucleon interaction as a function of the mass of the dark matter particle. This is the first search for dark matter particles produced in association with a Higgs boson decaying to two τ leptons.[Figure not available: see fulltext.]. © 2018, The Author(s).

SciVal Topic Prominence

Topic: jets | production | parton shower

Prominence percentile: 99.875

Author keywords

Dark matter Hadron-Hadron scattering (experiments)

Funding details

Funding sponsor	Funding number
California Earthquake Authority	

Metrics

0	Citations in Scopus
0	Field-Weighted Citation Impact

PlumX Metrics

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

Related research data

Search for dark matter produced in association with a Higgs boson decaying to $\gamma\gamma$ or $\tau^+\tau^-$ at $\sqrt{s}=13$ TeV

Ovcharova, Ana, et al
Deutsches Elektronen-Synchrotron, DESY, Hamburg

Search for dark matter produced in association with a Higgs boson decaying to $\gamma\gamma$ or $\tau^+\tau^-$ at $\sqrt{s}=13$ TeV

Ovcharova, Ana, et al
RWTH Aachen University

Acronym

Data linking provided by CEA

Related documents

Funding sponsor	Funding number	
Secretaría de Educación Superior, Ciencia, Tecnología e Innovación		Search for an exotic decay of the Higgs boson to a pair of light pseudoscalars in the final state with two b quarks and two T leptons in proton–proton collisions at $\sqrt{s}=13\text{TeV}$ SENESCYT
Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro		Sirunyan, A.M. , Tumasyan, A. , Adam, W. (2018) <i>Physics Letters, Section B: Nuclear, Elementary Particle and High-Energy Physics</i> FAPERJ
State Fund for Fundamental Research of Ukraine	Ukraine	Search for dark matter in association with a Higgs boson decaying to two photons at $\sqrt{s}=13\text{TeV}$ with the ATLAS detector SEER
CS Fund	Croatia	Aaboud, M. , Aad, G. , Abbott, B. (2017) <i>Physical Review D</i> CSF
Fuel Cell Technologies Program		Search for associated production of dark matter with a Higgs boson decaying to $b\bar{b}$ or $\gamma\gamma$ at $\sqrt{s}=13\text{TeV}$ FCT
Joint Institute for Nuclear Research		Sirunyan, A.M. , Tumasyan, A. , Adam, W. JINR (2017) <i>Journal of High Energy Physics</i>
Ministry of Education - Singapore		View all related documents based on references MOE
Pakistan Atomic Energy Commission		Find more related documents in Scopus based on: PAEC
Consejo Nacional de Ciencia y Tecnología		Authors > Keywords > CONACYT
Ministry for Business Innovation and Employment		MBIE
National Science and Technology Development Agency		NSTDA
Institute for Research in Fundamental Sciences		IPM
Missouri University of Science and Technology	Taipei	MST
Federación Española de Enfermedades Raras		FEDER
Hispanics in Philanthropy		HIP
Deutsche Forschungsgemeinschaft See opportunities by DFG		DFG
Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul		FAPERGS